



Wooden breast, white striping and spaghetti meat: chemical composition, technological quality, microbiological profile and sensory attributes of broiler breasts

Marija Bošković Cabrol^{a*}, Massimiliano Petracci^b and Angela Trocino^a

^a University of Padova, Department of Agronomy, Food, Natural Resources, Animal and Environment (DAFNAE), Viale dell'Università 16, 35020 Legnaro, Padova, Italy

^b Alma Mater Studiorum — University of Bologna, Department of Agricultural and Food Sciences, Cesena, Italy

ARTICLE INFO

Keywords:
Myopathies
P. major
Fatty acids
Texture
Cooking loss
Sensory profile

ABSTRACT

Poultry meat is consumed worldwide and its production is expected to increase in the upcoming years. Genetic selection in poultry focused on growth rate, feed conversion, and breast yield, resulting in the occurrence of white striping and wooden breast abnormalities and, most recently, spaghetti meat. These myopathies affect the quality traits of raw meat, including pH, color, water holding capacity, and cooking loss, which limit its further processing and decrease consumer acceptance. Additionally, the effects of myopathies on the chemical composition, i.e., reduced protein and essential amino acid content and increased fat contents, impair to some extent the nutritional value of the meat.

1. Introduction

Fast growth of poultry production with estimated increase of 16% by 2029 (FAO, 2020) is owed to the higher need for protein following rapid population growth (UN, 2019). The selection for fast weight gain puts excessive pressure on the *Pectoralis major* (*P. major*) muscle, resulting in high breast yield but also in a high occurrence of myopathies such as white striping (WS), wooden breast (WB) and spaghetti meat (SM) (Baldi et al., 2021; Petracci et al., 2019; Trocino et al., 2015; Kuttappan et al., 2013). On WS muscles, the presence of the white lines of intramuscular fat deposits can be seen (Kuttappan et al., 2013), whereas WB is characterized by hard-

ening of the muscle, and in severe cases, prominent ridge-like bulge, edema and/or scattered petechiae on the breast surface (Sihvo et al., 2014). *P. major* muscle with SM shows an overall impaired integrity with the separation of the fiber bundles composing the muscle tissue itself, resembling the appearance of spaghetti (Baldi et al., 2018). Although myopathies do not pose a threat to meat shelf life and microbiological safety (Gratta et al., 2019), they impair meat quality to some extent, affecting technological characteristics of raw meat and limiting their further processing (Gratta et al., 2019; Baldi et al., 2018; Mudalal et al., 2015; Trocino et al., 2015; Petracci et al., 2013). Despite carcasses affected by myopathies meeting the market quality standards (Baéza et

*Corresponding author: Marija Bošković Cabrol, marija.boskoviccabrol@unipd.it

Paper received May 1st 2023. Paper accepted May 12th 2023.

Published by Institute of Meat Hygiene and Technology — Belgrade, Serbia

This is an open access article under CC BY licence (<http://creativecommons.org/licenses/by/4.0>)

al., 2022) for weight and yield, muscle abnormalities impair the visual aspect of the meat, preventing the commercialization of whole breasts or, in cases of severe structural changes, requiring the breasts to be discarded as a waste (Zanetti et al., 2018).

2. Chemical composition of abnormal breast meat

Pathological alterations related to myopathies result in changes in the chemical composition of muscle tissue. Higher water and fat contents and lower ash and protein contents were reported in breasts affected by WS, WB, WB/WS and SM compared to their normal counterpart (Dalle Zotte et al., 2020; Gratta et al., 2019; Cai et al., 2018; Soglia et al., 2016 a, b; Zambonelli et al., 2016). The greater water content of WB and WS breasts is likely a consequence of inflammatory processes leading to fluid accumulation (Thanatsang et al., 2020; Sihvo et al., 2014). Indeed, Gratta et al., (2019) found higher moisture content in WB than in normal meat during the first 24 h *post-mortem*, while the differences among groups disappeared due to protein denaturation during 7 and 11 days of storage.

The reduction of protein content in WS, WB, and WB/WS meats due to fibrosis has been attributed to the replacement of muscle fibers with connective tissue and the aforementioned accumulation of adipose tissue (Dalle Zotte et al., 2020; Baldi et al., 2018; Soglia et al., 2016 a,b; Zambonelli et al., 2016), where a relationship between protein reduction and the degree of WS (Petracci et al., 2015; Kuttappan et al., 2013) has been also outlined. The protein decrease compared to normal breasts is followed by a reduction in essential amino acids in WS/WB and WB breast (Dalle Zotte et al., 2020; Thanatsang et al., 2020), which could seriously impair the meat nutritional value. The few studies about SM breasts found a reduction of protein content up to 10% compared to unaffected muscles (Baldi et al., 2018; 2019). Some authors found meat affected by SM, WB, WS/WB and WS/SM to display an increased collagen content compared to normal muscles, while others did not observe any difference (Baldi et al., 2018; Cai et al., 2018; Soglia et al., 2016 a,b). The degradation in the nutritional value of WS meat is also due to a lower digestibility and quality of amino acids in collagen compared to myofibrillar proteins (Huang and Ahn, 2018; Mudalal et al., 2014).

Compared with normal meat, the decrease in ash content in breast muscles associated with WS (Soglia

et al., 2018b; Mudalal et al., 2014), WB (Gratta et al., 2019; Cai et al., 2018; Soglia et al., 2016 a,b), and WS/WB (Dalle Zotte et al., 2020; Zambonelli et al., 2016) is probably due to pathological changes at cellular levels. After the initial degeneration, the damage to the sarcoplasmic reticulum surrounding muscle fibers increases Ca influx and activates Ca-dependent protease, triggering myofibers necrosis (Huang and Ahn, 2018). Increased Fe levels in WB could be due to hemorrhagic lesions (Thanatsang et al., 2020). In fact, Thanatsang et al., (2020) reported lower P and K and higher contents of Al, Ca, Fe, Na, and S in WB, and elevated Ca and Na content in SM compared to unaffected breasts. Zambonelli et al. (2016) found increased Ca and Na and decreased Mg and P, whereas Tasoniero et al., (2016) reported increased Fe and Na and reduced K and P contents in WS breasts.

Higher lipid content in WS and WB in comparison to normal meat are attributed to myodegeneration and replacement of degenerated muscle fibers by adipose tissue (Soglia et al., 2016a), often accompanied by changes in fatty acid profile. Baldi et al. (2018, 2019) reported a decrease in EPA and DHA in meat affected by SM compared to normal breasts, likely due to a different expression of the genes encoding for $\Delta 5$ and $\Delta 6$ desaturases (Soglia et al., 2016a). Gratta et al. (2019) reported chicken breasts affected by WB and WS display lower SFA and MUFA and higher PUFA due to an increase of both n-3 and n-6 FA compared to normal breasts. Lower SFA in WS and WB meat compared to the normal breasts was also reported in other studies (Soglia et al., 2016a; Kuttappan et al., 2012a). The mechanism behind changes in the FA profile of chicken meat affected by abnormalities is not completely clear. Possible explanations are related to the presence of inflammatory cells with a relatively high proportion of n-6 arachidonic acid in their membrane phospholipids (Calder, 2012) or the tendency of fiber type switching from fast, glycolytic IIB fibers, in which lipids are generally more saturated, to those within slow and oxidative fibers (Realini et al., 2013). Contrarily, Traffano-Schiffo et al. (2017) found that superficial meat layers affected by WS had higher SFA and MUFA and lower PUFA compared to normal meat. In fact, while most of the aforementioned studies reported changes in the overall *P. major*, some research found proximate composition changes only within the surface of the breast muscle (Baldi et al. 2019; Traffano-Schiffo et al., 2017). The chemical composition likely depends on the degree of structural changes: severe myopa-

thies will alter breast chemical composition at more profound parts of the muscle, whereas mild changes could only affect superficial meat layers.

Regarding oxidative stability, *Soglia et al.* (2016b) reported higher protein and lipid oxidation in WB and WS/WB compared to normal breasts, although no differences in total SFA, and PUFA were found between the former and the latter. The higher susceptibility of WB meat to protein oxidation was confirmed by *Thanatsang et al.* (2020). Despite higher lipid content, WS breasts do not seem to be more susceptible to lipid oxidation than unaffected ones (*Alnahhas et al.*, 2016; *Soglia et al.*, 2016b).

3. Technological quality of abnormal meat

In the presence of myopathies, reduction in protein content and degeneration of muscle fibers and myofibrils that usually entrap the majority of intracellular water are responsible for lower water holding capacity (WHC), marinade uptake and retention, emulsifying and gelling properties compared to normal breasts (*Petracci et al.*, 2019; *Xing et al.*, 2017; *Bowker and Zhuang*, 2016; *Soglia et al.*, 2016b). Whereas most of the findings on how WB affects WHC and drip and cooking losses are consistent (*Gratta et al.*, 2019; *Dalle Zotte et al.*, 2017; *Kuttappan et al.*, 2017a; *Trocino et al.*, 2015), the results on WS impact vary between studies. *Kuttappan et al.* (2017a) reported a decrease in drip loss in severe WB and WS/WB, but without differences between breasts with severe WS and normal ones as observed by *Kuttappan et al.* (2013) for cooking losses. Differently, *Petracci et al.* (2013) and *Tijare et al.* (2016) found an increase in cooking losses in raw breasts with severe WS. Because of the decreased total protein content but increased collagen and fat contents in WS breasts, their WHC, protein solubility and marinade uptake are lower than the normal muscles (*Sihvo et al.*, 2014). Regarding SM, this myopathy alone, or in combination with WS/WB, significantly increases drip losses (*Wang et al.*, 2023; *Tasoniero et al.*, 2020; *Baldi et al.*, 2018).

In terms of pH, some studies did not find changes in ultimate pH of muscle affected by myopathies (*Pascual et al.*, 2021; *Baldi et al.* 2018; *Trocino et al.*, 2015), while others reported elevated pH in WS, WB and SM breasts compared to their normal counterpart (*Wang et al.*, 2023; *Bordignon et al.*, 2022; *Tasoniero et al.*, 2016; *Mudalal et al.*, 2015; *Dalle Zotte et al.*, 2014; *Petracci et al.*, 2013). A proteomic study (*Kuttappan et al.*, 2017b) sug-

gested down-regulation of carbohydrate metabolic pathways related to reduced glycolysis, gluconeogenesis, TCA (tricarboxylic acid) cycle, glycogen degradation and pyruvate fermentation to lactate to be responsible for the reduced glycolytic potential and higher ultimate pH in affected breasts.

Regarding the effect of myopathies on breast color, inconsistent results have been reported. The meta-study of *Bordignon et al.* (2022) did not report an impact of WS, WB, or SM presence on breast color indices, in agreement with *Zambonelli et al.*, (2016) on WB and WS/WB breasts. Contrarily, *Mazzoni et al.* (2015) found a decrease in red and yellow indexes of WS breasts, whereas *Kuttappan et al.* (2017) reported an increase in yellowness in both WB and WS breasts when compared to control ones. Further, *Dalle Zotte et al.* (2017) noted elevated lightness in WB meat. Some authors suggested higher alteration in breast color as myopathy evolves to a severe degree (*Campo et al.*, 2020; *Tasoniero et al.*, 2016). *Campo et al.* (2020) observed a tendency to increase in lightness and yellowness in severe WB compared to unaffected meat, whereas the presence of WS also resulted in lighter, less red, and more yellow breasts. The presence of moderate SM resulted in higher yellowness. The higher yellowness of defective breasts arises from the elevated fat content (*Kuttappan et al.*, 2012a; 2013; *Petracci et al.*, 2015) and likely a high accumulation of dietary liposoluble pigments (*Campo et al.*, 2020).

Although the changes in texture of breasts affected by myopathies are documented, results differ between studies due to sample preparation (raw vs. cooked, surface layers vs. whole muscle), methods and parameters used (*Pascual et al.*, 2021; *Baldi et al.*, 2019; *Soglia et al.*, 2016a). The WB are often characterized by increased hardness of the muscle likely due to the higher collagen content of WB filets (*Baldi et al.*, 2019). However, the fibrosis in WB and WS/WB may not necessarily affect the texture properties of the breast, which was confirmed in several studies (*Xing et al.*, 2020; *Maxwell et al.*, 2018; *Sihvo et al.*, 2017; *Mudalal et al.*, 2015). Contrarily, SM and WS breasts usually display lower shear force than normal breasts (*Bordignon et al.*, 2022; *Baldi et al.*, 2019; *Tasoniero et al.*, 2016). Compared to normal breasts, lower hardness can be explained due to the higher fat content in WS breasts (*Radaelli et al.*, 2017; *Soglia et al.*, 2016a; *Mudalal et al.*, 2015) and to the reduced collagen cross-linking degree in SM ones (*Baldi et al.*, 2019, 2021). However, *Baldi et al.* (2019) did not find differences in the texture

between raw SM and normal samples, whereas cooking decreased hardness in SM. Pascual et al. (2021) reported only the Meullenet-Owens razor blade test to be able to detect the difference between SM and normal cooked breasts.

4. Microbiological quality of abnormal meat

To date, only a few studies investigated the microbiological status of meat with myopathies, suggesting it to be safe for human consumption. Gratta et al. (2019) reported higher initial total viable counts (TVC) and *Pseudomonas* spp. counts in normal breasts compared to WS and WB ones. Furthermore, normal breasts had the shortest TVC and *Pseudomonas* spp. Lag phase compared to WS and WB meat, resulting in the later microbial spoilage of breasts-affected abnormalities (5 vs. 6 days). *Enterobacteriaceae* spp. and lactic acid bacteria counts were also highest in unaffected breasts, followed by WS, and lowest in WB meat. Pereira et al. (2022) found lower *Staphylococcus* spp. and higher coliform counts in WS breasts, although bacterial counts neither in normal nor defective meat exceeded permitted limits after 12 months of freezing.

5. Sensory attributes of breasts with myopathies

As aforementioned, the impaired visual appearance and quality results in lower consumer acceptance and purchase intention of meat affected by myopathies (De Carvalho et al., 2020; Petracci et al., 2015). Consumers' acceptance of meat with myopathies strongly depends on the degree of muscle tissue changes. Kuttappan et al. (2012b) reported a decrease in consumer acceptance along with the severity of white striping, while Xing et al. (2020) found moderate and severe raw WB fillets to affect sensory acceptance due to their undesirable appearance, texture and drip loss, but found no differences between normal and mild WB. Indeed, the visual appearance of raw meat leads to meat rejection,

which is why some sensory studies are performed on cooked meat. Still, Tasoniero et al. (2016) reported more intensive off-odors in cooked WS fillets than in the normal ones, while López et al. (2022) observed a decrease in the cohesiveness and increase in the juiciness of severe WB. Although, in most of the studies, a deterioration of eating parameters is evident, de Almeida Assunção et al. (2020) found an increase in succulence in WB, probably due to a higher water content, to be well accepted by untrained panelists and increased final satisfaction with this meat. Furthermore, De Carvalho et al. (2020) reported lower acceptability of raw WS breast than unaffected counterparts, while upon cooking, consumers did not note differences between WS and normal chicken breasts for color, flavor, and overall acceptability but WS meat achieved higher scores for odor and texture compared to the normal breasts. Some efforts are made to eliminate or decrease the changes related to myopathies by marinating meat. However, Maxwell et al. (2018) reported differences in cooked meat sensory texture attributes related to WB to be lessened but not eliminated by vacuum-tumbling marination. Similarly, Jarvis et al. (2020) found neither traditional nor clean-label marinades to mask eating characteristics of severe WB, but, to some extent, a positive effect of marination was observed on mild WB. As far as SM goes, the effect of this muscle abnormality on sensory characteristics of breast meat has not yet been investigated.

6. Conclusion

Although the results between studies are sometimes inconsistent regarding how the different myopathies affect *P. major* in broilers, all studies reported at least one or multiple quality traits to be changed compared to unaffected breasts. Some underlying mechanisms behind changes in chemical compositions and quality of breasts affected by myopathies, especially less known SM, are not completely clear, emphasizing the need for further investigation on a molecular level.

Disclosure statement: No potential conflict of interest was reported by the authors.

Funding: Dr. Marija Bošković Cabrol acknowledges the European Union's Horizon 2020 research and innovation programme for the research grant she received under the Marie Skłodowska-Curie grant agreement No 101063055 (DeMYO project).

References

- Alnahhas, N., Berri, C., Chabault, M., Chartrin, P., Boulay, M., Bourin, M. C. & Bihan-Duval, L. (2016). Genetic parameters of white striping in relation to body weight, carcass composition, and meat quality traits in two broiler lines divergently selected for the ultimate pH of the pectoralis major muscle. *BMC genetics*, 17, 1–9, <https://doi.org/10.1186/s12863-016-0369-2>
- Baéza, E., Guillier, L. & Petracci, M. (2022). Production factors affecting poultry carcass and meat quality attributes. *Animal*, 16, e100331, <https://doi.org/10.1016/j.animal.2021.100331>
- Baldi, G., Soglia, F. & Petracci, M. (2021). Spaghetti meat abnormality in broilers: current understanding and future research directions. *Frontiers in Physiology*, 12, 684497, <https://doi.org/10.3389/fphys.2021.684497>
- Baldi, G., Soglia, F., Laghi, L., Tappi, S., Rocculi, P., Tavanliello, S., Prioriello, D., Mucci, R., Maiorano, G. & Petracci, M. (2019). Comparison of quality traits among breast meat affected by current muscle abnormalities. *Food Research International*, 115, 369–376, <https://doi.org/10.1016/j.foodres.2018.11.020>
- Baldi, G., Soglia, F., Mazzoni, M., Sirri, F., Canonico, L., Babini, E., Laghi, L., Cavani, C. & Petracci, M. (2018). Implications of white striping and spaghetti meat abnormalities on meat quality and histological features in broilers. *Animal*, 12, 164–173, <https://doi.org/10.1017/S1751731117001069>
- Bordignon, F., Xiccato, G., Boskovic Cabrol, M., Birolo, M. & Trocino, A. (2022). Factors affecting breast myopathies in broiler chickens and quality of defective meat: A meta-analysis. *Frontiers in Physiology*, e1322, <https://doi.org/10.3389/fphys.2022.933235>
- Bowker, B. & Zhuang, H. (2016). Impact of white striping on functionality attributes of broiler breast meat. *Poultry Science*, 95, 1957–1965, <https://doi.org/10.3382/ps/pew115>
- Cai, K., Shao, W., Chen, X., Campbell, Y. L., Nair, M. N., Suman, S. P., Beach, C. M., Guyton, M. C. & Schilling, M. W. (2018). Meat quality traits and proteome profile of woody broiler breast (pectoralis major) meat. *Poultry Science*, 97, 337–346, <https://doi.org/10.3382/ps/pex284>
- Calder, P. C. (2012). The role of marine omega-3 (n-3) fatty acids in inflammatory processes, atherosclerosis and plaque stability. *Molecular Nutrition & Food Research*, 56, 1073–1080, <https://doi.org/10.1002/mnfr.201100710>
- Campo, M. D. M., Mur, L., Guerrero, A., Barahona, M., Resconi, V. C., Magalhaes, D. R., Lisbinski, E., Boito, B., de Oliveira I. M. & Olleta, J. L. (2020). Differentiating breast myopathies through color and texture analyses in broiler. *Foods*, 9, 824, <https://doi.org/10.3390/foods9060824>
- Dalle Zotte, A., Cecchinato, M., Quartesan, A., Bradanovic, J., Tasoniero, G. & Puolanne, E. (2014). How does “wooden breast” myodegeneration affect poultry meat quality. In 60th International Congress of Meat, pp. 476–479.
- Dalle Zotte, A., Ricci, R., Cullere, M., Serva, L., Tenti, S. & Marchesini, G. (2020). Research Note: Effect of chicken genotype and white striping–wooden breast condition on breast meat proximate composition and amino acid profile. *Poultry Science*, 99, 1797–1803, <https://doi.org/10.1016/j.psj.2019.10.066>
- Dalle Zotte, A., Tasoniero, G., Puolanne, E., Remignon, H., Cecchinato, M., Catelli, E. & Cullere, M. (2017). Effect of “wooden breast” appearance on poultry meat quality, histological traits, and lesions characterization. *Czech Journal of Animal Science*, 62, 51–57, <https://doi.org/10.17221/54/2016-CJAS>
- de Almeida Assunção, A. S., Garcia, R. G., Komiyama, C. M., de Sena Gandra, É. R., de Souza, J. R., Dos Santos, W., Caldara, F.R. & Martins, R. A. (2020). Wooden breast myopathy on broiler breast fillets affects quality and consumer preference. *Tropical Animal Health and Production*, 52, 3555–3565, <https://doi.org/10.1007/s11250-020-02392-6>
- De Carvalho, L. M., Ventanas, S., Olegario, L. S., Madruga, M. S. & Estévez, M. (2020). Consumers awareness of white-striping as a chicken breast myopathy affects their purchasing decision and emotional responses. *LWT*, 131, e109809, <https://doi.org/10.1016/j.lwt.2020.109809>
- FAO (2020). Dairy and dairy products. *OECD-FAO Agricultural Outlook*, 2029, 174–183.
- Gratta, F., Fasolato, L., Birolo, M., Zomeño, C., Novelletti, E., Petracci, M., Pascual, A., Xiccato G. & Trocino, A. (2019). Effect of breast myopathies on quality and microbial shelf life of broiler meat. *Poultry Science*, 98, 2641–2651, <https://doi.org/10.3382/ps/pez001>
- Huang, X. & Ahn, D. U. (2018). The incidence of muscle abnormalities in broiler breast meat—a review. *Korean Journal for Food Science of Animal Resources*, 38, 835, <https://doi.org/10.5851/kosfa.2018.e2>
- Jarvis, T., Byron, M., Von Staden, M., Crist, C., Zhang, X., Rowe, C., Smith, B. & Schilling, M. W. (2020). Quality differences in wooden and normal broiler breast meat marinated with traditional and clean label marinades. *Meat and Muscle Biology*, 4, 1–13, <https://doi.org/10.22175/mmb.9458>
- Kuttappan, V. A., Brewer, V. B., Apple, J. K., Waldroup, P. W. & Owens, C. M. (2012a). Influence of growth rate on the occurrence of white striping in broiler breast fillets. *Poultry Science*, 91, 2677–2685, <https://doi.org/10.3382/ps.2012-02259>
- Kuttappan, V. A., Lee, Y. S., Erf, G. F., Meullenet, J. F., McKee, S. R. & Owens, C. M. (2012b). Consumer acceptance of visual appearance of broiler breast meat with varying degrees of white striping. *Poultry Science*, 91, 1240–1247, <https://doi.org/10.3382/ps.2011-01947>
- Kuttappan, V. A., Shivaprasad, H. L., Shaw, D. P., Valentine, B. A., Hargis, B. M., Clark, F. D., McKee, R.S. & Owens, C. M. (2013). Pathological changes associated with white striping in broiler breast muscles. *Poultry Science*, 92, 331–338, <https://doi.org/10.3382/ps.2012-02646>
- Kuttappan, V. A., Owens, C. M., Coon, C., Hargis, B. M. & Vazquez-Anon, M. (2017a). Incidence of broiler breast myopathies at 2 different ages and its impact on selected raw meat quality parameters. *Poultry Science*, 96, 3005–3009, <https://doi.org/10.3382/ps/pex072>
- Kuttappan, V. A., Bottje, W., Ramnathan, R., Hartson, S. D., Coon, C. N., Kong, B. W., Owens, M.C., Vazquez-Añon,

- M. & Hargis, B. M. (2017b). Proteomic analysis reveals changes in carbohydrate and protein metabolism associated with broiler breast myopathy. *Poultry Science*, 96, 2992–2999, <https://doi.org/10.3382/ps/pex069>
- Maxwell, A. D., Bowker, B. C., Zhuang, H., Chatterjee, D. & Adhikari, K. (2018). Descriptive sensory analysis of marinated and non-marinated wooden breast fillet portions. *Poultry Science*, 97, 2971–2978, <https://doi.org/10.3382/ps/pey145>
- Mazzoni, M., Petracci, M., Meluzzi, A., Cavani, C., Clavanzani, P. & Sirri, F. (2015). Relationship between pectoralis major muscle histology and quality traits of chicken meat. *Poultry Science*, 94, 123–130, <https://doi.org/10.3382/ps/peu043>
- Mudalal, S., Babini, E., Cavani, C. & Petracci, M. (2014). Quantity and functionality of protein fractions in chicken breast fillets affected by white striping. *Poultry Science*, 93, 2108–2116, <https://doi.org/10.3382/ps.2014-03911>
- Mudalal, S., Lorenzi, M., Soglia, F., Cavani, C. & Petracci, M. (2015). Implications of white striping and wooden breast abnormalities on quality traits of raw and marinated chicken meat. *Animal*, 9, 728–734, <https://doi.org/10.1017/S175173111400295X>
- Pascual, A. G., Trocino, A., Susta, L. & Barbut, S. (2021). Comparing three textural measurements of chicken breast fillets affected by severe wooden breast and spaghetti meat. *Italian Journal of Animal Science*, 20, 465–471, <https://doi.org/10.1080/1828051X.2021.1893134>
- Pereira, M. R., Mello, J. L., Oliveira, R. F., Villegas-Caylahua, E. A., Cavalcanti, E. N., Fidelis, H.X.A., Ferrari, B. F., Giampietro-Ganeco, A., Souza, A. P. & Borba, H. (2022). Effect of freezing on the quality of breast meat from broilers affected by White Striping myopathy. *Poultry Science*, 101, 101607, <https://doi.org/10.1016/j.psj.2021.101607>
- Petracci, M., Mudalal, S., Bonfiglio, A. & Cavani, C. (2013). Occurrence of white striping under commercial conditions and its impact on breast meat quality in broiler chickens. *Poultry Science*, 92, 1670–1675, <https://doi.org/10.3382/ps.2012-03001>
- Petracci, M., Mudalal, S., Soglia, F. & Cavani, C. (2015). Meat quality in fast-growing broiler chickens. *World's Poultry Science Journal*, 71, 363–374, <https://doi.org/10.1017/S0043933915000367>
- Petracci, M., Soglia, F., Madruga, M., Carvalho, L., Ida, E. & Estévez, M. (2019). Wooden-breast, white striping, and spaghetti meat: causes, consequences and consumer perception of emerging broiler meat abnormalities. *Comprehensive Reviews in Food Science and Food Safety*, 18, 565–583, <https://doi.org/10.1111/1541-4337.12431>
- Radaelli, G., Piccirillo, A., Birolo, M., Bertotto, D., Gratta, F., Ballarin, C., Vascellari, M., Xiccato, G. & Trocino, A. (2017). Effect of age on the occurrence of muscle fiber degeneration associated with myopathies in broiler chickens submitted to feed restriction. *Poultry Science*, 96, 309–319, <https://doi.org/10.3382/ps/pew270>
- Realini, C. E., Vénien, A., Gou, P., Gatellier, P., Perez-Juan, M., Danon, J. & Astruc, T. (2013). Characterization of *Longissimus thoracis*, *Semitendinosus* and *Masseter* muscles and relationships with technological quality in pigs. 1. Microscopic analysis of muscles. *Meat Science*, 94, 408–416, <https://doi.org/10.1016/j.meatsci.2013.03.009>
- Sihvo, H. K., Immonen, K. & Puolanne, E. (2014). Myodegeneration with fibrosis and regeneration in the pectoralis major muscle of broilers. *Veterinary Pathology*, 51, 619–623, <https://doi.org/10.1177/0300985813497488>
- Sihvo, H. K., Lindén, J., Airas, N., Immonen, K., Valaja, J. & Puolanne, E. (2017). Wooden breast myodegeneration of pectoralis major muscle over the growth period in broilers. *Veterinary Pathology*, 54, 119–128, <https://doi.org/10.1177/03009858166580>
- Soglia, F., Laghi, L., Canonico, L., Cavani, C. & Petracci, M. (2016b). Functional property issues in broiler breast meat related to emerging muscle abnormalities. *Food Research International*, 89, 1071–1076, <https://doi.org/10.1016/j.foodres.2016.04.042>
- Soglia, F., Mudalal, S., Babini, E., Di Nunzio, M., Mazzoni, M., Sirri, F., Cavani, C. & Petracci, M. (2016a). Histology, composition, and quality traits of chicken Pectoralis major muscle affected by wooden breast abnormality. *Poultry Science*, 95, 651–659, <https://doi.org/10.3382/ps/pev353>
- Tasoniero, G., Cullere, M., Cecchinato, M., Puolanne, E. & Dalle Zotte, A. (2016). Technological quality, mineral profile, and sensory attributes of broiler chicken breasts affected by White Striping and Wooden Breast myopathies. *Poultry Science*, 95, 2707–2714, <https://doi.org/10.3382/ps/pew215>
- Thanatsang, K. V., Malila, Y., Arayamethakorn, S., Srimarut, Y., Tatiyaborworntam, N., Uengwetwanit, T., Panya, A., Rungrassamee, W. & Visessanguan, W. (2020). Nutritional properties and oxidative indices of broiler breast meat affected by wooden breast abnormality. *Animals*, 10, e2272, <https://doi.org/10.3390/ani10122272>
- Tijare, V. V., Yang, F. L., Kuttappan, V. A., Alvarado, C. Z., Coon, C. N. & Owens, C. M. (2016). Meat quality of broiler breast fillets with white striping and woody breast muscle myopathies. *Poultry Science*, 95, 2167–2173, <https://doi.org/10.3382/ps/pew129>
- Traffano-Schiffo, M. V., Castro-Giraldez, M., Colom, R. J. & Fito, P. J. (2017). Development of a spectrophotometric system to detect white striping physiopathy in whole chicken carcasses. *Sensors*, 17, e1024, <https://doi.org/10.3390/s17051024>
- Trocino, A., Piccirillo, A., Birolo, M., Radaelli, G., Bertotto, D., Filiou, E., Petracci, M. & Xiccato, G. (2015). Effect of genotype, gender and feed restriction on growth, meat quality and the occurrence of white striping and wooden breast in broiler chickens. *Poultry Science*, 94, e2996–3004, <https://doi.org/10.3382/ps/pev296>
- UN (United Nations), (2019). World population prospects 2019 — highlights. New York, pp 46. (<https://www.un.org/development/desa/publications/world-population-prospects-2019-highlights.html>)
- Wang, C., Che, S., Susta, L. & Barbut, S. (2023). Textural and physical properties of breast fillets with myopathies (wooden breast, white striping, spaghetti meat) in Canadian fast-growing broiler chickens. *Poultry Science*, 102, 102309, <https://doi.org/10.1016/j.psj.2022.102309>
- Xing, T., Zhao, X., Cai, L., Guanghong, Z. & Xu, X. (2017). Effect of salt content on gelation of normal and wooden breast myopathy chicken pectoralis major meat batters. *International Journal of Food Science & Technology*, 52, 2068–2077, <https://doi.org/10.1111/ijfs.13485>

Xing, T., Zhao, X., Zhang, L., Li, J. L., Zhou, G. H., Xu, X. L. & Gao, F. (2020). Characteristics and incidence of broiler chicken wooden breast meat under commercial conditions in China. *Poultry Science*, 99, 620–628, <https://doi.org/10.3382/ps/pez560>

Zambonelli, P., Zappaterra, M., Soglia, F., Petracci, M., Sirri, F., Cavani, C. & Davoli, R. (2016). Detection of differentially expressed genes in broiler pectoralis major muscle affected by White Striping–Wooden Breast

myopathies. *Poultry Science*, 95, 2771–2785, <https://doi.org/10.3382/ps/pew268>

Zanetti, M. A., Tedesco, D. C., Schneider, T., Teixeira, S. T. F., Daroit, L., Pilotto, F., Dickel, E.L., Santos, P.S. & dos Santos, L. R. (2018). Economic losses associated with wooden breast and white striping in broilers. *Semina: Ciências Agrárias*, 39, 887–891, <https://doi.org/10.5433/1679-0359.2018v39n2p887>