



# The influence of different gas mixtures on the shelf life of fresh beef

Milan Milijašević<sup>a\*</sup>, Jelena Babić Milijašević<sup>a</sup>, Ivan Nastasijević<sup>a</sup> and Slavica Vesković Moračanin<sup>a</sup>

<sup>a</sup> Institute of Meat Hygiene and Technology, Kačanskog 13, 11000 Belgrade, Serbia

## ARTICLE INFO

### Keywords:

Modified atmosphere  
Beef round primal cuts  
Microbiological status  
Gas mixture

## ABSTRACT

The paper presents the results of testing the influence of different gas mixtures on the shelf life of different beef round primal cuts. A total of five gas mixtures were used, and the change in microbiological status (total number of bacteria, *Escherichia coli*, *Salmonella* spp., *Listeria monocytogenes*, *Proteus* spp., sulphite-reducing clostridia), change in pH and sensory characteristics of the meat were monitored. The results of this study showed that beef packaged in a gas mixture consisting of 70% O<sub>2</sub> and 30% CO<sub>2</sub> was the most acceptable in terms of sensory characteristics and microbiological status.

## 1. Introduction

Nowadays, food must be healthy, minimally processed and attractively packaged, as consumer expectations are constantly growing (Martinez *et al.*, 2006). As a consequence, the demands to be met by manufacturers of food and packaging materials have also increased (McMillin *et al.*, 1999; Brody, 2003). Today's consumers are also very sensitive when it comes to the use of additives in the food industry. Public demands to be able to get fresh food at any time are increasing, and food safety and easy availability of all kinds of foods are very important. That is why it is becoming increasingly difficult to meet consumer expectations. It is clear that time, as a factor in food production, plays an important role. From the moment when the fruit is harvested, the cattle are slaughtered, or the fish is caught, the race against time begins. From that moment on, natural degradation and deterioration threaten the quality and sustainability of the product. However, external factors such as the hygiene of the production process, temperature, etc., also pose dangers to the usability of

the product. Therefore, the ways in which the food will be handled during the production, packaging, or cold chain processes are very important. Special emphasis should be placed on the packaging stage, because the packaging method can enable greater sustainability of the product (Phillips *et al.*, 2001).

Over the past two decades, modified atmosphere packaging (MAP) has become the dominant form of packaging in the meat industry (Robertson, 1993). The main reasons that stimulated the development of this technology are the constant increase in the consumption of fresh meat, the increase in the population in cities and the reduction of natural food sources. MAP can be defined as the removal of air from the package and its replacement with a specific gas or mixture of gases. The purpose of this technology is to extend the sustainability of food by preventing or slowing biochemical processes (oxidation of fats, formation of metmyoglobin), the growth of spoilage bacteria and the degree of product respiration. Numerous literature data indicate that several gases, such as carbon dioxide (CO<sub>2</sub>), nitro-

\*Corresponding author: Milan Milijašević, [milan.milijasevic@inmes.rs](mailto:milan.milijasevic@inmes.rs)

Paper received July 22<sup>nd</sup> 2023. Paper accepted May 31<sup>st</sup> 2023.

Published by Institute of Meat Hygiene and Technology — Belgrade, Serbia

This is an open access article under CC BY licence (<http://creativecommons.org/licenses/by/4.0>)

gen ( $N_2$ ), oxygen ( $O_2$ ) and carbon monoxide (CO), are used individually or in different combinations in MAP technology (Xiong, 1999).  $CO_2$  shows antimicrobial activity, mostly against Gram-negative bacteria and psychrotrophic pathogens such as *Yersinia enterocolitica* and *Aeromonas hydrophila* (Brody, 1989), but this gas does not significantly inhibit the growth of *Listeria monocytogenes*.  $CO_2$  also does not affect the growth of *Clostridium botulinum* (Sinell, 1988).  $O_2$  is used to preserve the bright red color of the meat and prevent the growth of anaerobic bacteria (Martinez et al., 2006).  $N_2$  is an inert gas that replaces  $O_2$  in the mixture and prolongs the product's shelf life, and also prevents rancidity and the growth of aerobic bacteria (Blakistone, 1998).  $N_2$ , due to its low solubility, has a role to prevent the collapse of the package. CO is very effective in preserving the red color of fresh meat, as it has a 20-fold greater affinity for binding to myoglobin, compared to  $O_2$  (Boeckman, 2006).

The aim of this experiment was to investigate the effect of different gas mixtures on the color, pH value and microbiological status of beef packaged in a modified atmosphere.

## 2. Materials and methods

The study was carried out in a medium-sized meat processing plant. Beef round parts were used as the matrix. The average temperature of the pieces of

meat during packaging was  $3.1^\circ C$ , and the pH was 5.75. Twenty-four hours after packaging, the pieces of meat were cut into smaller pieces, where the average temperature of the meat was  $4.2^\circ C$ , and the pH was 5.54. Temperature and pH values were measured every three days using a Testo 205 apparatus (Testo AG, Germany). Pieces of meat, 1–2 cm thick, were placed in Cryovac LidSys polystyrene containers with a depth of 50 mm (Sealed Air, USA), in which there were papers for absorbing meat juice (Sealed Air, USA). The pieces of meat were then packed in a modified protective atmospheres and stored for 28 days at  $4^\circ C$ . A total of five different gas mixtures (Messer AG, Germany) were used in the study (Table 1).


The microbiological status and pH were examined 2, 6, 13, 17 and 28 days after setting up the study. The color of the meat was evaluated organoleptically, at the beginning and at the end of the study, without instrumental measurement. Microbiological tests were performed according to validated ISO methods. These tests were the total number of viable aerobic bacteria (TVA), *Escherichia coli*, *Salmonella* spp., *Listeria monocytogenes*, *Proteus* spp. and sulphite-reducing clostridia.

The change in the total number of bacteria is presented in Figure 1.


In cuts of meat packed in a gas mixture consisting of 30%  $CO_2$  and 70%  $N_2$ , and in 100%  $CO_2$ ,

**Table 1.** Composition of gas mixtures used in the modified atmosphere packaging for beef.

	Mix 1	Mix 2	Mix 3	Mix 4	Mix 5
$CO_2$	30%	100%	50%	30%	20%
$O_2$	-	-	-	70%	20%
$N_2$	70%	-	50%	-	60%



**Figure 1.** Total viable aerobic counts measured in beef packed in different gas mixtures (see Table 1) during the study.



**Figure 2.** pH measured in beef packed in different gas mixtures (see Table 1) during the study.

the total number of bacteria remained low almost throughout the study. In beef pieces packed in gas mixture containing 20% and 70% O<sub>2</sub>, during the entire study, there was a gradual increase in TVA, with a sudden increase between days 13 and 17.

The presence of pathogenic bacteria during the entire experiment (*E. coli*, *Salmonella* spp., *L. monocytogenes*, sulphite-reducing clostridia) was not recorded in any sample of beef packed in MAP.

The results of monitoring the pH change are shown in Figure 2. A significant increase in the pH value occurred in beef packaged in a gas mixture containing 70% O<sub>2</sub>. Other mixtures showed a slow but steady increase in pH. A significant increase in pH value was noted around day 15, for all gas mixtures.

Pieces of beef packaged in gas mixtures containing a high percentage of CO<sub>2</sub> acquired a dark green color, and meat packaged in a mixture consisting of 30% CO<sub>2</sub> and 70% O<sub>2</sub> retained the most acceptable red color.

### 3. Discussion and results

Increasing the concentration of oxygen from 20% to 70% in the packaging has a positive effect on the color stability of the micro-fabricated parts of the beef leg. These results were also confirmed in research by other authors (Seyfert *et al.*, 2007; Djenane *et al.*,

2001; Brody, 1989; Martinez *et al.*, 2006). This is probably due to the formation of a larger amount of oxymyoglobin in the meat in proportion to the higher concentration of oxygen in the gas mixture. Pieces of beef packed in mixtures containing a high percentage of CO<sub>2</sub> acquired an unacceptable dark green color due to the formation of metmyoglobin.

Beef that was packed in gas mixtures with a higher percentage of CO<sub>2</sub> showed lower pH values. This is explained by the dissolution of CO<sub>2</sub> in the aqueous phase of the meat, whereby carbonic acid is formed, which leads to a decrease in pH. These statements were made by other authors in their works (Phillips, 1996; Antoniewski *et al.*, 2007; Arnold *et al.*, 1993; Bendall *et al.*, 1972; Klettner, 2004; Hess *et al.*, 1980).

The TVA was higher in beef packed in the gas mixture with a higher (70%) O<sub>2</sub> than in the 20% O<sub>2</sub> mixture. This was due to the aerobic conditions that prevailed in the packaging and the slightly higher pH of the beef in the high-O<sub>2</sub> mixture that favoured the development of microorganisms.

### 4. Conclusion

The TVA was higher in beef packed in a mixture with a higher percentage of O<sub>2</sub>, due to the aerobic conditions that prevailed in the packaging and slightly higher pH values that favoured the development of microorganisms.

**Disclosure statement:** No potential conflict of interest was reported by the authors.

**Funding:** This study was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia, according to the provisions of the Contract on research financing in 2023 (No 451-03-47/2023-01/200050 dated 03.02.2023).

## References

- Antoniewski, N., Barringer, A., Knipe, L. & Zerby, N., (2007).** Effect of a gelatin coating on the shelf life of fresh meat. *Journal of Food Science*, 72(6), E382–E387.
- Arnold, N., Arp, C., Scheller, K., Williams, N. & Schaefer, M., (1993).** Tissue equilibrium and subcellular distribution of vitamin E relative to myoglobin and lipid oxidation in displayed beef. *Journal of Animal Science*, 71, 105–118.
- Bendall, R. & Taylor, A., (1972).** Consumption of oxygen by the muscles of beef animals and related species. II. Consumption of oxygen by post-rigor muscles. *Journal of the Science of Food and Agriculture*, 23, 707–719.
- Blakistone, A., (1998).** Introduction. In B. A. Blakistone (Ed.), Principles and applications of modified atmosphere packaging of foods (pp. 1–13). Gaithersburg, Maryland: Aspen Publishers.
- Boeckman, M., (2006).** Regulatory status of carbon monoxide for meat packaging. In Proceedings 59<sup>th</sup> reciprocal meat conference (pp. 59–60), 18–21 June 2006, Champaign-Urbana, Illinois, USA.
- Brody, A. L. (2003).** “Nano, Nano” Food Packaging Technology. *Journal of Food Technology*, 12, 52–54.
- Brody, L., (1989).** Modified atmosphere/vacuum packaging of meat. In A. L. Brody (Ed.), Controlled/modified atmosphere/vacuum packaging of foods (pp. 17–37). Trumbull, Connecticut: Food & Nutrition Press.
- Djenane, D., Sánchez-Escalante, A., Beltrán, A. & Roncalés, P., (2001).** Extension of the Retail Display Life of Fresh Beef Packaged in Modified Atmosphere by Varying Lighting Conditions. *Journal of Food Science*, 66(1), 181–185.
- Hess, E., Ruosch, W. & Breer, C., (1980).** Extending the shelf-life of prepacked fresh meat. *Fleischwirtschaft*, 60 (8), 1513–1516.
- Klettner, P.-G., (2004).** Schutzgasverpackung bei Frischfleisch (Rückenmuskulatur von Schwein und Rind) unter besonderer Berücksichtigung von Kohlenmonoxid, Mitteilungsblatt der Fleischforschung, Kulmbach 43. Nr. 166, 331–342.
- Martínez, L., Djenane, D., Cilla, I., Beltrán, A. & Roncalés, P., (2006).** Effect of varying oxygen concentrations on the shelf-life of fresh pork sausages packaged in modified atmosphere. *Food Chemistry*, 94, 219–225.
- McMillin, W., Huang, Y., Ho, P. & Smith, S., (1999).** Quality and shelf-life of meat in case-ready modified atmosphere packaging. In Y. L. Xiong, F. Shahidi, & C. T. Ho (Eds.), Quality attributes in muscle foods (pp. 73–93). New York: ACS Symposium Series, Plenum Publishing Corporation.
- Phillips, C., (1996).** Modified atmosphere packaging and its effects on the microbiological quality and safety of produce. *International Journal of Food Science and Technology*, 31, 463–479.
- Phillips, L., Faustman, C., Lynch, P., Govoni, E., Hoagland, A. & Zinn, A., (2001).** Effect of dietary a-tocopherol supplementation on color and lipid stability in pork. *Meat Science*, 58, 389–393.
- Robertson, G., (1993).** Food packaging principles and practice. New York: Marcel Dekker 431–469.
- Seyfert, M., Mancini, A., Hunt, C., Tang, J. & Faustman, C., (2007).** Influence of carbon monoxide in package atmospheres containing oxygen on colour, reducing activity, and oxygen consumption of five bovine muscles. *Meat Science*, 75, 432–442.
- Sinell, H.-J., (1988).** Mikrobiologie des Fleisches. In Prändl-Fischer-Schnidhofer-Sinell, 1988, Fleisch, Ulmer, 1988, Stuttgart.
- Xiong, Li, (1999).** Modified Atmosphere Packaging. Dept. of Food Science, Penn State University.